Jumping into Statistics: Introduction to Study Design and Statistical Analysis for Medical Research Using JMP Pro Statistical Software

WINTER/SPRING 2021

DR. CYNDI GARVAN & DR. TERRIE VASILOPOULOS

Meet the Instructors



CYNTHIA GARVAN, MA, PHD

Research Professor in Anesthesiology cgarvan@anest.ufl.edu



TERRIE VASILOPOULOS, PHD

Research Assistant Professor in Anesthesiology and Orthopaedics and Rehabilitation

tvasilopoulos@anest.ufl.edu

Course Objectives

- Review fundamentals of study design and research methodology
- Understand how to choose best statistical test for your research question
- Practice basic statistical analysis use JMP Pro Software

Course Topics

- Asking a Good Research Question
- Life Cycle of Research and Scientific Method
- Study Design
- Data types and Database Construction
- Descriptive Statistics
- Data Visualization
- Population and Sample,Probability, Statistical Inference

- How to Chose Correct Statistical Method and Run Some Analyses
 - T-tests, ANOVA, Non-Parametric
 - Chi-square, odds ratio, relative risk
 - Regression and Correlation
 - Survival Analysis
 - Test Diagnostics (e.g. sensitivity, specificity, etc.)
- Comparing Statistical Modeling and Machine Learning

Study Design: Sample Size and Power Calculations

2/10/2021

What is statistical power?

- Ability of a study to detect differences or effects if they exist.
- What influences power?
 - Effect size (e.g. differences, correlation)
 - Sample Variation (e.g. SD)
 - Sample size
 - In study planning, we aim to determine sample size needed in order adequately power a study

Understanding Errors

Your Statistical Decision	True state of null hypothesis	
	H ₀ True (example: the drug doesn't work)	H ₀ False (example: the drug works)
$\begin{array}{c} \text{Reject } H_0 \\ \text{(ex: you conclude that the drug works)} \end{array}$	Type I error (a) False positive Incorrectly rejecting null	Correct
Do not reject H_0 (ex: you conclude that there is insufficient evidence that the drug works)	Correct	Type II Error (β) False Negative Incorrectly accepting null

For most research, α is set to 0.05...i.e. willing to accept, at most, a 5% chance of obtaining a false positive

That's why we look for p < 0.05 in statistical testing

Determining statistical power

- Power = 1 beta (Type II error rate)
 - Probability of rejecting null when it is actually false
 - i.e. statistically significant result when an effect exists
 - Typical Threshold = 80%
 - Note: Type I and Type II error are inversely related

Determining statistical power

- What information do we need for sample size determination?
 - Power threshold (80%)
 - Smallest effect size of interest
 - Mean difference
 - Proportion difference/Event rate difference
 - Correlation
 - Odds Ratio
 - **■** Type I error (alpha = 0.05)
 - Variability
 - SD
 - Baseline proportions, event rates in controls
 - Best if these values are informed by previous work!

Power Programs

- •JMP Pro (but it is limited)
- G*Power
 - More options (correlations, regression, odds ratio, repeated measures)
 - https://www.psychologie.hhu.de/arbeitsgruppen/allgemeine-psychologie-undarbeitspsychologie/gpower.html

GLIMMPSE

- Mixed Models
- https://glimmpse.samplesizeshop.org/
- Power and Sample Size
 - http://powerandsamplesize.com/
- Open Epi
 - Note: has not been updated in a while, but links and programs still work
 - https://www.openepi.com/Menu/OE Menu.htm
- •For more complex designs, should seek consult from a statistician

Demo!

Questions?

JMP Pro!

https://software.ufl.edu/

